



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Baird's Sandpiper near Washington, D. C.—On Sept. 3, 1894, while collecting Sandpipers and other birds on the floating confervæ on the flats in the Potomac River, off the mouth of Four Mile Run, Alexandria County, Virginia, I took a fine immature male specimen of *Tringa bairdii*. This specimen has been identified by Mr. Ridgway and is the first recorded instance from the Virginias and the District of Columbia. It was alone at the time, though hundreds of individuals of *Ereunetes pusillus*, *Tringa maculata* and *Totanus flavipes* were flying about or feeding on the surface of the confervæ.—R. S. MATTHEWS, *Washington, D. C.*

The Ruff and Western Sandpiper near Washington, D. C.—With other Waders on the confervæ off the mouth of Four Mile Run, Alexandria County, Virginia, I collected on September 3, 1894, an immature female *pavoncella pugnax*. The condition of the specimen would warrant the belief that it was bred on this continent.

I also took on September 8, at the same place, an immature male *Ereunetes occidentalis*, which is the first record from this locality. In contrast with the excessive fatness of many specimens of the Least and other Sandpipers taken lately at the same place, both the above, especially the Sandpiper, were quite lean. A second specimen, an immature female, was taken on Sept. 11, at the same place, by Mr. R. S. Matthews. These specimens were seen and identified by Mr. R. Ridgway.—WILLIAM PALMER, *Washington, D. C.*

An Asiatic Cuckoo on the Pribylof Islands, Alaska.—An adult male Cuckoo (No. 118864 U. S. N. M. Coll.) that I collected among the sand dunes of Northeast Point, St. Paul's Island, on July 4, 1890, has been identified by Dr. L. Stejneger as *Cuculus canorus telephonus* (Heine). This bird, whose summer habitat is Japan and Kamtschatka, has several times been taken on Bering Island. When collected it was busily engaged capturing some large flies, which are abundant on these islands, and with which its stomach was literally packed. It had been seen by the natives in the same place for more than two weeks, and was probably the same individual seen by myself on June 13, when becalmed in a fog off the eastern side of the same island, on which occasion it circled overhead like a Gull for some time while calmly inspecting the boat, and then moved off northwards.—WILLIAM PALMER, *Washington, D. C.*

The Capture of *Basilinna leucotis* in Southern Arizona.—While collecting in the high Chiricahua Mountains, in southern Arizona during the past summer, the writer had the pleasure of securing a specimen of *Basilinna leucotis*, a new bird to the United States.

During the early part of June a camp was made at Fly Park, a well wooded area southeast of the head of Pinery Cañon, at an altitude of about 10,000 feet. A boreal honeysuckle (*Lonicera involucrata*) grows